



TITLE:

CONSTRUCTION OF FOLD MAP OF LENS SPACE $L(p,1)$ WHERE SINGULAR SET IS A TORUS (Singularity theory of differential maps and its applications)

AUTHOR(S):

Yamamoto, Minoru

CITATION:

Yamamoto, Minoru. CONSTRUCTION OF FOLD MAP OF LENS SPACE $L(p,1)$ WHERE SINGULAR SET IS A TORUS
(Singularity theory of differential maps and its applications). 数理解析研究所講究録 2017, 2049: 45-56

ISSUE DATE:

2017-10

URL:

<http://hdl.handle.net/2433/237051>

RIGHT:

CONSTRUCTION OF FOLD MAP OF LENS SPACE $L(p, 1)$ WHERE SINGULAR SET IS A TORUS

弘前大学教育学部 山本稔
MINORU YAMAMOTO

FACULTY OF EDUCATION, HIROSAKI UNIVERSITY

1. INTRODUCTION

Throughout the report, all manifolds and maps are differentiable of class C^∞ . Let $f : M \rightarrow \mathbb{R}^p$ be a map of a closed n -dimensional manifold M into \mathbb{R}^p ($n \geq p$). We denote by $S(f)$ the set of points in M where the rank of the differential of f is strictly less than p . We say that $S(f) \subset M$ is a *singular set* of f and $f(S(f)) \subset \mathbb{R}^p$ is a *contour* of f .

Let $f : M \rightarrow \mathbb{R}^3$ be a map of a closed connected oriented 3-dimensional manifold M into \mathbb{R}^3 . For any $q \in S(f)$ of $f : M \rightarrow \mathbb{R}^3$, if we can choose local coordinates (u_1, u_2, u_3) centered at q and (v_1, v_2, v_3) centered at $f(q)$ respectively such that f has the following form:

$$(1.1) \quad (v_1 \circ f, v_2 \circ f, v_3 \circ f) = (u_1, u_2, u_3^2),$$

then we call f a *fold map*. It is known that if $f : M \rightarrow \mathbb{R}^3$ is a fold map, then $S(f)$ is a closed orientable surface (not necessary connected) and $f|S(f) : S(f) \rightarrow \mathbb{R}^3$ is an immersion. If $f|S(f)$ is an immersion with normal crossings, we call f a *stable fold map*.

Eliashberg [2] showed that if a closed surface V splits M into two manifolds M_1, M_2 with $\partial M_1 = \partial M_2 = V$, then there exists a fold map $f : M \rightarrow \mathbb{R}^3$ such that $S(f) = V$. Here, M_1 and M_2 are not necessary connected. In this report, we apply Eliashberg's theorem to a lens space $L(p, 1)$ and construct a stable fold map $f : M \rightarrow \mathbb{R}^3$ such that $S(f) = T^2$ is a Heegaard surface of $L(p, 1)$ ($p \geq 2$).

The author would like to thank the organizers Professor Masatomo Takahashi and Professor Takahiro Yamamoto for organizing and including him in the conference, "Singularity theory of differential maps and its applications". He also would like to thank Professor Kenta Hayano, Professor Goo Ishikawa, Professor Yusuke Mizota, Professor Takashi Nishimura, Professor Osamu Saeki, Professor Kentaro Saji and Professor Kazuto Takao for invaluable comments at his talk of this conference.

2. DESCRIPTION OF A STABLE FOLD MAP

In this section, we explain a method to depict a stable fold map $f : M \rightarrow \mathbb{R}^3$. In the following, we assume that M is a closed connected oriented 3-dimensional manifold and that \mathbb{R}^3 and \mathbb{R}^2 are oriented.

The author has been supported by Grant-in-Aid for Young Scientists (B) JSPS (No. 25800032).

For a stable fold map $f : M \rightarrow \mathbb{R}^3$ such that $S(f) = V$ and $M = M_1 \cup_V M_2$, we remark that $f|_{M_1}$ and $f|_{M_2}$ are immersions and extensions of $f|_V$. We assume that $f|_{M_1}$ is an orientation preserving immersion and $f|_{M_2}$ is an orientation reversing immersion. The orientation on M_1 induces the orientation on V as follows. For $q \in V$, let $\{n_1, n_2, nt_3\}$ be the basis of $T_q(M_1)$ which defines the orientation on M_1 and n_1 the outward normal vector. Then the orientation on $V = \partial(M_1)$ is defined by $\{n_2, n_3\}$.

By Bruce and Kirk's theorem [1], there exists an orthogonal projection $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ such that $\pi \circ f|_V : V \rightarrow \mathbb{R}^2$ is a stable map. It is well-known that a stable map satisfies the following properties.

Proposition 2.1 ([3]). *A smooth map $g : N \rightarrow \mathbb{R}^2$ of a closed surface N into \mathbb{R}^2 is a stable map if and only if the following conditions are satisfied.*

- (1) For every $q \in S(g)$, there exist local coordinates (u_1, u_2) and (v_1, v_2) around q and $g(q)$ respectively such that one of the following holds:
 - (i) $(v_1 \circ g, v_2 \circ g) = (u_1, u_2^2)$, q : fold point,
 - (ii) $(v_1 \circ g, v_2 \circ g) = (u_1, u_2^3 - u_1 u_2)$, q : cusp point.
- (2) If q is a cusp point of g , then $g^{-1}(g(q)) \cap S(g) = \{q\}$,
- (3) $g|_{S(g) \setminus \{\text{set of cusp points of } g\}}$ is an immersion with normal crossings.

In the following, we set $f_V^\pi = \pi \circ f|_V$. Let $q \in V$ be a cusp point of a stable map $f_V^\pi : V \rightarrow \mathbb{R}^2$. For a sufficiently small neighborhood U of $f_V^\pi(q)$, the map $f_V^\pi|_{U'} : U' \rightarrow U$ has degree ± 1 , where U' is the component of $(f_V^\pi)^{-1}(U)$ containing q . If the degree of q is $+1$ (resp. -1), then we should paint q and $f_V^\pi(q)$ red (resp. blue).

For each $t \in \mathbb{R}$, a plane $\{(t, y, z) \in \mathbb{R}^3 \mid y, z \in \mathbb{R}\}$ is denoted by \mathbb{R}_t^2 . Then, for almost all $t \in \mathbb{R}$, $f(V) \cap \mathbb{R}_t^2$ consists of immersed circles (or an empty set), $f(M_i) \cap \mathbb{R}_t^2$ consists of immersed surfaces (or an empty set) and $f(M_i) \cap \mathbb{R}_t^2$ is an extension of $f(V) \cap \mathbb{R}_t^2$. Therefore, from the pictures $f(M_1) \cap \mathbb{R}_{t_1}^2, f(M_1) \cap \mathbb{R}_{t_2}^2, \dots, f(M_1) \cap \mathbb{R}_{t_n}^2$ and $f(M_2) \cap \mathbb{R}_{t_1}^2, f(M_2) \cap \mathbb{R}_{t_2}^2, \dots, f(M_2) \cap \mathbb{R}_{t_n}^2$, we can see the immersed 3-dimensional manifold $f(M_1), f(M_2)$ and the image of the stable fold map $f(M)$. Note that the planes $\mathbb{R}_{t_1}^2, \mathbb{R}_{t_2}^2, \dots, \mathbb{R}_{t_n}^2$ can be chosen from the picture of the contour $f_V^\pi(S(f_V^\pi)) \subset \mathbb{R}^2$.

For a fold point $q \in S(f_V^\pi)$ of f_V^π , there exist local coordinates (u_1, u_2, u_3) and (v_1, v_2) around $q \in M$ and $\pi \circ f(q) \in \mathbb{R}^2$ such that

$$(v_1 \circ (\pi \circ f), v_2 \circ (\pi \circ f)) = (u_1, u_2^2 \pm u_3^2)$$

holds. Here, $S(f)$ corresponds to $\{u_3 = 0\}$. If q corresponds to the map $(v_1 \circ (\pi \circ f), v_2 \circ (\pi \circ f)) = (u_1, u_2^2 + u_3^2)$ (resp. $(v_1 \circ (\pi \circ f), v_2 \circ (\pi \circ f)) = (u_1, u_2^2 - u_3^2)$), then we should paint q and $\pi \circ f(q)$ red (resp. blue). From the local picture around $S(f_V^\pi)$, we have the following.

- On each connected component of $S(f_V^\pi) \setminus \{\text{cusp points}\}$, it should be colored by red or blue.
- If two connected components of $S(f_V^\pi) \setminus \{\text{cusp points}\}$ adjacent to the same cusp point, then they are painted by the different colors. See Figure 2 of the web version for example.

3. CONSTRUCTION OF A STABLE FOLD MAP $f^{(2,1)} : L(2, 1) \rightarrow \mathbb{R}^3$

In this section, we construct a stable fold map $f^{(2,1)} : L(2, 1) \rightarrow \mathbb{R}^3$ such that $S(f^{(2,1)}) = T^2$ is a Heegaard surface of $L(2, 1)$.

(Step 1.) Let $g : V \rightarrow \mathbb{R}^2$ be a stable map of a closed connected surface V to \mathbb{R}^2 such that the contour $g(S(g))$ and the inverse images $g^{-1}(\mathbb{R}_{t_1}) \cap V, \dots, g^{-1}(\mathbb{R}_{t_{11}}) \cap V$ are depicted in Figure 1. Here, \mathbb{R}_t is a line defined by $\mathbb{R}_t = \{(t, y) \in \mathbb{R}^2 | y \in \mathbb{R}\}$.

Since $g^{-1}(\mathbb{R}_{t_1}) \cap V, \dots, g^{-1}(\mathbb{R}_{t_{11}}) \cap V$ can be seen as a sequence of immersed curves in $\mathbb{R}_{t_i}^2$, we can lift the stable map $g : V \rightarrow \mathbb{R}^2$ to a generic immersion $g' : V \rightarrow \mathbb{R}^3$ such that $g = \pi \circ g'$. From Figure 1, we can check that V is a torus. In the following, we consider that the sequence in Figure 1 is the sequence of immersed circles $g'(V) \cap \mathbb{R}_{t_1}^2, \dots, g'(V) \cap \mathbb{R}_{t_{11}}^2$.

(Step 2.) From Figure 1, we construct two kinds of sequences of immersed surfaces which are extensions of immersed circles $g'(V) \cap \mathbb{R}_{t_1}^2, \dots, g'(V) \cap \mathbb{R}_{t_{11}}^2$. Figure 2 represents one sequence of immersed surfaces and Figure 3 represents another sequence.

By combining the immersed surfaces in Figure 2, we have an immersion $f_1 : M_1 \rightarrow \mathbb{R}^3$ which is one extension of the generic immersion $g' : V \rightarrow \mathbb{R}^3$. Also, by combining the immersed surfaces in Figure 3, we have an immersion $f_2 : M_2 \rightarrow \mathbb{R}^3$ which is another extension of the generic immersion $g' : V \rightarrow \mathbb{R}^3$. We define the orientation of M_1 (resp. M_2) so as the immersion f_1 (resp. f_2) is an orientation preserving (resp. orientation reversing). In Figure 2 (resp. Figure 3), green bands explain how each immersed surface $f_1(M_1) \cap \mathbb{R}_{t_i}^2$ (resp. $f_2(M_2) \cap \mathbb{R}_{t_i}^2$) is obtained as the extension of the immersed circles $g'(V) \cap \mathbb{R}_{t_i}^2$. See the web version.

(Step 3) Let $C \subset V$ be a circle such that $C \subset S(g)$ and the image $g(C)$ is depicted as gray thick lines in Figure 4. By a regular homotopy of f_2 , we can check that M_2 is a solid torus and C is a meridian circle of M_2 . By a regular homotopy of f_1 , we can check that M_1 is a solid torus and C is a $(2, 1)$ -curve of M_1 . That is, C turns twice in the longitude direction and once in the meridian direction on M_1 . Therefore, by attaching these immersions f_1 and f_2 , we obtain a stable fold map $f^{(2,1)} = f_1 \cup f_2 : M_1 \cup_V M_2 = L(2, 1) \rightarrow \mathbb{R}^3$ such that $S(f^{(2,1)}) = V = T^2$ is a Heegaard surface.

4. CONSTRUCTION OF A STABLE FOLD MAP $f^{(p,1)} : L(p, 1) \rightarrow \mathbb{R}^3$

In this section, we construct a stable fold map $f^{(p,1)} : L(p, 1) \rightarrow \mathbb{R}^3$ such that $S(f^{(p,1)}) = T^2$ is a Heegaard surface of $L(p, 1)$ ($p \geq 2$).

(Step 1.) Let $g' : V \rightarrow \mathbb{R}^3$ be a generic immersion of a closed connected surface V to \mathbb{R}^3 such that $g = \pi \circ g'$ is a stable map and the contour $g(S(g))$ is depicted in Figure 5. Let U be a subset of \mathbb{R}^2 depicted in Figure 5. The image $g(V) \cap (\mathbb{R}^2 \setminus U)$ of Figure 5 is the same as that of Figure 1. Therefore, in Figure 6, we only describe a sequence of immersed arcs $g'(V) \cap \pi^{-1}(\mathbb{R}_t \cap U)$. From Figures 5 and 6, we can check that V is a torus.

(Step 2.) From Figure 6, we construct two kinds of sequences of immersed surfaces which are extensions of immersed arcs $g'(V) \cap \pi^{-1}(\mathbb{R}_t \cap U)$. Figure 7 represents one sequence of immersed surfaces and Figures 8 represents another sequence. By combining the immersed surfaces in Figure 7, we have




FIGURE 1. The contour of $g : V \rightarrow \mathbb{R}^2$ and the sequence of sectional faces of $g(V)$ or $g'(V)$.

an immersion $f_1 : M_1 \rightarrow \mathbb{R}^3$ which is one extension of the generic immersion $g' : V \rightarrow \mathbb{R}^3$. Also, by combining the immersed surfaces in Figure 8, we have an immersion $f_2 : M_2 \rightarrow \mathbb{R}^3$ which is another extension of the generic immersion $g' : V \rightarrow \mathbb{R}^3$. We define the orientation of M_1 (resp. M_2) so as the immersion f_1 (resp. f_2) is an orientation preserving (resp. orientation reversing). In Figure 7 (resp. Figure 8), green bands explain how each immersed




FIGURE 2. The sequence of sectional faces of $f_1(M_1)$.

surface $f_1(M_1) \cap \pi^{-1}(\mathbb{R}_t \cap U)$ (resp. $f_2(M_2) \cap \pi^{-1}(\mathbb{R}_t \cap U)$) is obtained as the extension of the immersed arcs $g'(V) \cap \pi^{-1}(\mathbb{R}_t \cap U)$. See the web version.

(Step 3) Let $C \subset V$ be a circle such that $C \subset S(g)$ and the image $g(C)$ is depicted as gray thick lines in Figure 9. By a regular homotopy of f_2 , we can check that M_2 is a solid torus and C is a meridian circle of M_2 . By a regular homotopy of f_1 , we can check that M_1 is a solid torus and C is a $(p, p-1)$ -curve of M_1 . Therefore, by attaching these immersions f_1 and




FIGURE 3. The sequence of sectional faces of $f_2(M_2)$.

f_2 , we obtain a stable fold map $f_1 \cup f_2 : M_1 \cup_V M_2 = L(p, p-1) \rightarrow \mathbb{R}^3$ such that $S(f_1 \cup f_2) = V = T^2$ is a Heegaard surface. Since $L(p, p-1)$ is diffeomorphic to $L(p, 1)$, $f^{(p,1)} = f_1 \cup f_2$ is a desired stable fold map.




FIGURE 4. The image of the curve C which is a meridian circle of M_2 .




FIGURE 5. The contour of $g : V \rightarrow \mathbb{R}^2$.




FIGURE 6. The sequence of the sectional faces of $g'(V)$.





FIGURE 7. The sequence of sectional faces of $f_1(M_1)$.

5. REMARKS AND PROBLEMS

In Sections 3 and 4, we only construct a stable fold map of $L(p, 1)$ whose singular set is a genus one Heegaard surface. Therefore, we have a following problem.

Problem 5.1. Construct a stable fold map $f^{(p,q)} : L(p, q) \rightarrow \mathbb{R}^3$ such that $S(f^{(p,q)})$ is a genus one Heegaard surface ($p - 1 > q > 1$).

For the stable fold map $f^{(2,1)} : L(2, 1) \rightarrow \mathbb{R}^3$ of Section 3, we can check that $(f^{(2,1)})^{-1}(f^{(2,1)}(L(2, 1)) \cap \mathbb{R}_{t_6}^2)$ is a torus in $L(2, 1)$. Let $\mathbb{R}_{(-\infty, t_6]}$ and

FIGURE 8. The sequence of sectional faces of $f_2(M_2)$.

$\mathbb{R}_{[t_6, \infty)}^3$ be half spaces defined by $\mathbb{R}_{(-\infty, t_6]}^3 = \{(x, y, z) \in \mathbb{R}^3 \mid x \in (-\infty, t_6]\}$ and $\mathbb{R}_{[t_6, \infty)}^3 = \{(x, y, z) \in \mathbb{R}^3 \mid x \in [t_6, \infty)\}$. Let N_1 and N_2 be submanifolds of $L(2, 1)$ defined by $N_1 = L(2, 1) \cap (f^{(2,1)})^{-1}(\mathbb{R}_{(-\infty, t_6]}^3)$ and $N_2 = L(2, 1) \cap (f^{(2,1)})^{-1}(\mathbb{R}_{[t_6, \infty)}^3)$. We have a following problem.

Problem 5.2. Does the decomposition $N_1 \cup_{T^2} N_2$ represent a genus one Heegaard splitting of $L(2, 1)$?

Let $S^3 = D_1^3 \cup_{S_1^2} S^2 \times I \cup_{S_2^2} D_2^3$ be a decomposition of S^3 and $e : S^3 \rightarrow \mathbb{R}^3$ be a stable fold map such that $S(e) = S_1^2 \cup S_2^2$ and $e|_{D_1^3}$ and $e|_{D_2^3}$ are orientation




FIGURE 9. The image of the curve C which is a meridian circle of M_2 .

preserving immersions and $e|S^2 \times I$ is an orientation reversing immersion. Figure 10 represents the contour of the stable map $e_{S_1^2 \cup S_2^2}^\pi : S_1^2 \cup S_2^2 \rightarrow \mathbb{R}^2$ and the sequence of the sectional faces of $e(S_1^2 \cup S_2^2)$. Figure 11 (resp. Figure 12) represents the sequence of the sectional faces of $e(D_1^3)$ (resp. $e(D_2^3)$) and Figure 13 represents the sequence of the sectional faces of $e(S^2 \times I)$.




FIGURE 10. The contour of $e_{S_1^2 \cup S_2^2}^\pi : S_1^2 \cup S_2^2 \rightarrow \mathbb{R}^2$ and the sequence of the sectional faces of $e(S_1^2 \cup S_2^2)$.

By a connected sum of the two stable fold maps $f^{(p,1)} \sharp e$ and the Eliashberg's trick which is introduced in [2], we have a stable fold map $f_2^{(p,1)} : L(p, 1) \rightarrow \mathbb{R}^3$ such that $S(f_2^{(p,1)}) = T^2 \sharp T^2$ is a genus two Heegaard surface ($p \geq 2$). The contour of $\pi \circ f_2^{(p,1)}|S(f_2^{(p,1)})$ is depicted in Figure 14. By repeating the above operation, we have a stable fold map $f_k^{(p,1)} : L(p, 1) \rightarrow \mathbb{R}^3$ such that $S(f_k^{(p,1)}) = \sharp^k T^2$ is a genus k Heegaard surface ($p \geq 2$).




FIGURE 11. The sequence of the sectional faces of $e(D_1^3)$.




FIGURE 12. The sequence of the sectional faces of $e(D_2^3)$.




FIGURE 13. The sequence of the sectional faces of $e(S^2 \times I)$.




FIGURE 14. The contour of $\pi \circ f_2^{(p,1)}|S(f_2^{(p,1)})$.

If we use the Eliashberg's trick for the stable fold map $e : S^3 \rightarrow \mathbb{R}^3$, we have a stable fold map $f^{(1,0)} : S^3 \rightarrow \mathbb{R}^3$ such that $S(f^{(1,0)}) = T^2$ is a genus one Heegaard surface. Therefore, we also have a stable fold map $f_k^{(1,0)} : S^3 \rightarrow \mathbb{R}^3$ such that $S(f_k^{(1,0)}) = \#^k T^2$ is a genus k Heegaard surface. We have a following problem.

Problem 5.3. Construct a nontrivial stable fold map $f : L(p, p-1) \rightarrow \mathbb{R}^3$ such that $S(f)$ is a genus k Heegaard surface ($p \geq 1, k \geq 2$).

Let $\text{SI}(3, 1)$ be the group of oriented bordism classes of immersions of closed oriented 3-dimensional manifolds in \mathbb{R}^4 and $\text{SFold}(3, 0)$ the group of oriented fold cobordism classes of fold maps of closed oriented 3-dimensional manifolds into \mathbb{R}^3 . Let $K : S^3 \rightarrow \mathbb{R}^4$ be an immersion which is constructed from the track of the standard Froissart-Morin's eversion $S^2 \times I \rightarrow \mathbb{R}^4$. Hughes [5] showed that the immersion K is a generator of $\text{SI}(3, 1)$. Hirato-Takase [4] showed that the homomorphism $\mathfrak{m} : \text{SFold}(3, 0) \rightarrow \text{SI}(3, 1)$ is an isomorphism. Since we can check that e and $f^{(1,0)} : S^3 \rightarrow \mathbb{R}^3$ are oriented fold cobordant, and that the bordism class of K is equal to $\mathfrak{m}(e)$, the stable fold map $f^{(1,0)} : S^3 \rightarrow \mathbb{R}^3$ is a generator of $\text{SFold}(3, 0)$. This also shows that $f^{(1,0)} : S^3 \rightarrow \mathbb{R}^3$ is a generator of the third stable stem π_3^S .

REFERENCES

- [1] J. W. Bruce and N. P. Kirk, *Generic projections of stable mappings*, Bull. London Math. Soc. **32** (2000), 718–728.
- [2] Y. Eliashberg, *On singularities of folding type*, Math. USSR Izvestija **4** (1970), 1119–1134.
- [3] M. Golubitsky and V. Guillemin, *Stable mappings and their singularities*, Graduate Texts in Mathematics, Vol. 14, Springer-Verlag, New York, Heidelberg, 1973.
- [4] Y. Hirato and M. Takase, *Compositions of equi-dimensional fold maps*, Fund. Math. **216** (2012), 119–128.
- [5] J. F. Hughes, *Bordism and regular homotopy of low-dimensional immersions*, Pacific J. Math. **156** (1992), 155–184.